

# The XENON1T Low-Energy Excess (And Other Exciting Results)

Dr. Erwann Masson

IJCLab – Orsay, France

On behalf of the XENON Collaboration

Particle Physics Seminars at BNL – 19 Nov. 2020



université  
PARIS-SACLAY

Xe  
XENON  
Dark Matter Project

# On Dark Matter & Liquid Xenon

# Dark Matter in a nutshell

- ▶ A **non-luminous matter** is needed to explain what is observed in the Universe **at all scales**

# Dark Matter in a nutshell

- A **non-luminous matter** is needed to explain what is observed in the Universe **at all scales** ( $1 \text{ pc} \approx 3.26 \text{ ly}$ )

20-60 kpc



NGC 6503, Hubble Space Telescope

# Dark Matter in a nutshell

- A **non-luminous matter** is needed to explain what is observed in the Universe **at all scales** ( $1 \text{ pc} \approx 3.26 \text{ ly}$ )

20-60 kpc





2-10 Mpc



# Dark Matter in a nutshell

- ▶ A **non-luminous matter** is needed to explain what is observed in the Universe **at all scales** ( $1 \text{ pc} \approx 3.26 \text{ ly}$ )
- ▶ Standard cosmological model → **27%** of non-baryonic, non-relativistic, and almost non-interacting matter

20-60 kpc  
2-10 Mpc  
4 Gpc



# Dark Matter in a nutshell

- ▶ A **non-luminous matter** is needed to explain what is observed in the Universe **at all scales** ( $1 \text{ pc} \approx 3.26 \text{ ly}$ )
- ▶ Standard cosmological model → **27%** of non-baryonic, non-relativistic, and almost non-interacting matter
- ▶ Most promising candidate in particle physics  
→ **Weakly Interacting Massive Particles (WIMPs,  $\chi$ )**



## WIMP (expected) ID card

- ▶ High mass,  $m_\chi \in \text{GeV}/c^2\text{--TeV}/c^2$
- ▶ Low velocity,  $v_0 \approx 220 \text{ km/s}$  (locally)
- ▶ Very low interaction cross section,  
 $\sigma_{\text{int}} \in 10^{-41}\text{--}10^{-48} \text{ cm}^2$
- ▶ Elastically scatters off atomic nuclei  
→ **Nuclear Recoils (NR)  $\lesssim 100 \text{ keV}$**





# Dark Matter in a nutshell

- ▶ A **non-luminous matter** is needed to explain what is observed in the Universe **at all scales** ( $1 \text{ pc} \approx 3.26 \text{ ly}$ )
- ▶ Standard cosmological model → **27%** of non-baryonic, non-relativistic, and almost non-interacting matter
- ▶ Most promising candidate in particle physics  
→ **Weakly Interacting Massive Particles (WIMPs,  $\chi$ )**





# Dark Matter in a nutshell

- ▶ A **non-luminous matter** is needed to explain what is observed in the Universe **at all scales** ( $1 \text{ pc} \approx 3.26 \text{ ly}$ )
- ▶ Standard cosmological model → **27%** of non-baryonic, non-relativistic, and almost non-interacting matter
- ▶ Most promising candidate in particle physics  
→ **Weakly Interacting Massive Particles (WIMPs,  $\chi$ )**



20-60 kpc  
2-10 Mpc  
4 Gpc



# The XENON Collaboration



# The XENON Collaboration (in COVID times)



XENON Technical Meeting, May 12-14, 2020


Andrii Terliuk (MPIK/Uni He...)


Alexey Elykov

Ethan Brown

Christopher Hils (JGU-Mai...)


Michele Iacovacci





**INFN**  
LNGS





**INFN**  
LNGS



Dual-Phase TPC

Water Čerenkov  
Muon Veto



Cryogenics & Xenon Purification

Electronics & Data Acquisition

Xenon Storage,  
Recovery & Distillation


# The XENON Program

PRL 100 (2008) 021303  
PRD 94 (2016) 122001  
PRL 121 (2018) 111302



**XENON10**  
2005–2007

25 kg LXe  
15 cm drift length  
 $\sigma_{\text{SI}} \sim 9 \times 10^{-44} \text{ cm}^2$   
at 100 GeV/c<sup>2</sup> (2007)



**XENON100**  
2009–2016

161 kg LXe  
30 cm drift length  
 $\sigma_{\text{SI}} \sim 10^{-45} \text{ cm}^2$   
at 50 GeV/c<sup>2</sup> (2016)



**XENON1T**  
2016–2018

3.2 t LXe  
1 m drift length  
 $\sigma_{\text{SI}} \sim 4 \times 10^{-47} \text{ cm}^2$   
at 30 GeV/c<sup>2</sup> (2018)


# The XENON Program

PRL 100 (2008) 021303  
PRD 94 (2016) 122001  
PRL 121 (2018) 111302



**XENON10**  
2005–2007

25 kg LXe  
15 cm drift length  
 $\sigma_{\text{SI}} \sim 9 \times 10^{-44} \text{ cm}^2$   
at  $100 \text{ GeV}/c^2$  (2007)



**XENON100**  
2009–2016

161 kg LXe  
30 cm drift length  
 $\sigma_{\text{SI}} \sim 10^{-45} \text{ cm}^2$   
at  $50 \text{ GeV}/c^2$  (2016)



**XENON1T**  
2016–2018

3.2 t LXe  
1 m drift length  
 $\sigma_{\text{SI}} \sim 4 \times 10^{-47} \text{ cm}^2$   
at  $30 \text{ GeV}/c^2$  (2018)



**XENONnT**  
2020–2025


NOW

8.4 t LXe  
1.5 m drift length  
 $\sigma_{\text{SI}} \sim 2 \times 10^{-48} \text{ cm}^2$   
at  $50 \text{ GeV}/c^2$  (20 t  $\times$  yr)

# Liquid Xenon as a detection medium

## Liquid Xenon (LXe) ID card

- Atomic number  $Z = 54$
- High mass number,  $\langle A \rangle = 131$
- High density at 177 K,  $\langle \rho \rangle = 2.86 \text{ g/cm}^3$
- No long-lived radioisotopes in WIMP ROI
- Efficient UV scintillator (178 nm)




PRC 81 (2010) 025808

# Liquid Xenon as a detection medium

## Liquid Xenon (LXe) ID card

- ▶ Atomic number  $Z = 54$
- ▶ **High mass number**,  $\langle A \rangle = 131$
- ▶ High density at 177 K,  $\langle \rho \rangle = 2.86 \text{ g/cm}^3$
- ▶ No long-lived radioisotopes in WIMP ROI
- ▶ Efficient UV scintillator (178 nm)

▶ **Maximised** interaction cross section ( $\propto A^2$ )




PRC 81 (2010) 025808

# Liquid Xenon as a detection medium

## Liquid Xenon (LXe) ID card

- ▶ Atomic number  $Z = 54$
- ▶ High mass number,  $\langle A \rangle = 131$
- ▶ **High density** at 177 K,  $\langle \rho \rangle = 2.86 \text{ g/cm}^3$
- ▶ No long-lived radioisotopes in WIMP ROI
- ▶ Efficient UV scintillator (178 nm)

- ▶ **Maximised** interaction cross section ( $\propto A^2$ )
- ▶ High stopping power and **self-shielding**




PRC 81 (2010) 025808

# Liquid Xenon as a detection medium

## Liquid Xenon (LXe) ID card

- ▶ Atomic number  $Z = 54$
- ▶ High mass number,  $\langle A \rangle = 131$
- ▶ High density at 177 K,  $\langle \rho \rangle = 2.86 \text{ g/cm}^3$
- ▶ **No long-lived radioisotopes** in WIMP ROI
- ▶ Efficient UV scintillator (178 nm)

- ▶ **Maximised** interaction cross section ( $\propto A^2$ )
- ▶ High stopping power and **self-shielding**
- ▶ **Free** from harmful intrinsic background




PRC 81 (2010) 025808

# Liquid Xenon as a detection medium

## Liquid Xenon (LXe) ID card

- Atomic number  $Z = 54$
- High mass number,  $\langle A \rangle = 131$
- High density at 177 K,  $\langle \rho \rangle = 2.86 \text{ g/cm}^3$
- No long-lived radioisotopes in WIMP ROI
- Efficient **UV scintillator** (178 nm)

- **Maximised** interaction cross section ( $\propto A^2$ )
- High stopping power and **self-shielding**
- **Free** from harmful intrinsic background
- Powerful DM discrimination combining **scintillation and ionisation** properties




PRC 81 (2010) 025808

# Liquid Xenon as a detection medium

## Liquid Xenon (LXe) ID card


- ▶ Atomic number  $Z = 54$
- ▶ High mass number,  $\langle A \rangle = 131$
- ▶ High density at 177 K,  $\langle \rho \rangle = 2.86 \text{ g/cm}^3$
- ▶ No long-lived radioisotopes in WIMP ROI
- ▶ Efficient **UV scintillator** (178 nm)

- ▶ **Maximised** interaction cross section ( $\propto A^2$ )
- ▶ High stopping power and **self-shielding**
- ▶ **Free** from harmful intrinsic background
- ▶ Powerful DM discrimination combining **scintillation and ionisation** properties
- ▶ **Scalable** → well suited to DM search and evolving projects like **XENON**




PRC 81 (2010) 025808


# Detecting particles with a dual-phase TPC




# Detecting particles with a dual-phase TPC




# Detecting particles with a dual-phase TPC




# Detecting particles with a dual-phase TPC





# Detecting particles with a dual-phase TPC






# Detecting particles with a dual-phase TPC



# Reconstructing energy






# Reconstructing energy




$$\begin{aligned} E &= W(n_\gamma + n_e) \\ &= W\left(\frac{S1}{g_1} + \frac{S2}{g_2}\right) \end{aligned}$$

# Reconstructing energy



$$E = W(n_\gamma + n_e)$$
$$= W \left( \frac{S1}{g_1} + \frac{S2}{g_2} \right)$$



# The evolution of LXe TPCs


Total LXe mass  
**3.2 tonnes**



XENON10



XENON100



LUX



PANDA X-II




XENON1T

Xe

2016

2000 kg

0.2



XENONnT

Xe

2020

6000 kg

0.02  
(Goal)



2005

2009

2013

2016

2020



ACTIVE LIQUID XENON  
TARGET MASS

22 kg

105 kg


250 kg

580 kg

6000 kg




LOW ENERGY  
ER BACKGROUND  
 $(t \times d \times \text{keV})^{-1}$



What XENON1T  
found so far

# High-mass WIMP-nucleus interactions


PRL 121 (2018) 111302



- A few events in the region of interest but **no significant excess** over background

# High-mass WIMP-nucleus interactions


PRL 121 (2018) 111302




- ▶ A few events in the region of interest but **no significant excess** over background
- ▶ To date, **strongest limit** to the WIMP-nucleon cross section for WIMPs **above  $6 \text{ GeV}/c^2$**
- ▶ Unprecedented minimum of  $4.1 \times 10^{-47} \text{ cm}^2$  at  $30 \text{ GeV}/c^2$  (90% C.L.)

# Light DM search (S2-only)

PRL 123 (2019) 251801



(A): SI DM-nucleus scattering




(C): SD DM-neutron scattering


- ▶ Analysis using the **ionisation signal (S2) only** → access to 2–3 times lower energies

# Light DM search (S2-only)

PRL 123 (2019) 251801



(A): SI DM-nucleus scattering




(C): SD DM-neutron scattering

- ▶ Analysis using the **ionisation signal (S2) only** → access to 2–3 times lower energies
- ▶ Best sensitivity to low-mass DM-matter interactions **below 6 GeV/c $^2$**

# Light DM search (S2-only)


PRL 123 (2019) 251801



- ▶ Analysis using the **ionisation signal (S2) only** → access to 2–3 times lower energies
- ▶ Best sensitivity to low-mass DM-matter interactions **below  $6 \text{ GeV}/c^2$**
- ▶ **New constraints on various models**, from DM-electron scatterings to axion-like particles

# Light DM search (Migdal & Brems)



PRL 123 (2019) 241803



- ▶ Taking advantage of the **post-NR lag** between the nucleus and orbital electrons motion  
→ **Migdal** and **Bremsstrahlung**  $\lesssim 1$  keV **ER** radiation

# Light DM search (Migdal & Brems)




PRL 123 (2019) 241803



- ▶ Taking advantage of the **post-NR lag** between the nucleus and orbital electrons motion
  - **Migdal** and **Bremsstrahlung**  $\lesssim 1 \text{ keV ER}$  radiation
- ▶ Using S2 alone → sensitivity to DM **down to 85 MeV/ $c^2$**

# Light DM search (Migdal & Brems)

PRL 123 (2019) 241803




- ▶ Taking advantage of the **post-NR lag** between the nucleus and orbital electrons motion  
→ **Migdal** and **Bremsstrahlung**  $\lesssim 1$  keV **ER** radiation
- ▶ Using S2 alone → sensitivity to DM **down to 85 MeV/c²**
- ▶ Best upper limits for DM masses **below  $\sim 2$  GeV/c²**


# Light DM search (Migdal & Brems)

PRL 123 (2019) 241803

- ▶ Taking advantage of nucleus and orbital motion
- Migdal and Brems scattering
- ▶ Using S2 alone → sensitivity improvement
- ▶ Best upper limits for DM mass




Leading (most of) the searched DM mass range!



# 2ν Double Electron Capture in $^{124}\text{Xe}$



Nature 568 (2019) 7753



- ▶ Extremely rare process, never observed before

# 2ν Double Electron Capture in $^{124}\text{Xe}$



Nature 568 (2019) 7753



- Extremely rare process, never observed before
- K-shell electron capture → X-rays and  $\text{e}_{\text{Auger}}$  (64.3 keV)

# 2v Double Electron Capture in $^{124}\text{Xe}$




Nature 568 (2019) 7753




- Extremely rare process, never observed before
- K-shell electron capture → **X-rays and  $\text{e}_{\text{Auger}}$  (64.3 keV)**
- $^{124}\text{Xe} \sim 1 \text{ kg/t LXe} \rightarrow \text{126 2vECEC events}$  ( $4.4\sigma$  above bkg.)

# 2v Double Electron Capture in $^{124}\text{Xe}$

Nature 568 (2019) 7753




- ▶ Extremely rare process, never observed before
- ▶ K-shell electron capture → **X-rays and  $e_{\text{Auger}}$  (64.3 keV)**
- ▶  $^{124}\text{Xe} \sim 1 \text{ kg/t LXe} \rightarrow \text{126 2vECEC events}$  ( $4.4\sigma$  above bkg.)
- ▶ Resulting half-life →  $T_{1/2} = (1.8 \pm 0.5_{\text{stat}} \pm 0.1_{\text{sys}}) \times 10^{22} \text{ yr}$
- ▶ Unprecedented sensitivity to **rare decay searches**



# 2v Double Electron Capture in $^{124}\text{Xe}$

Nature 568 (2019) 7753



# The Low-Energy ER Excess

# An excess of events

PRD 102 (2020) 072004

PHYSICAL REVIEW D 102, 072004 (2020)

Featured in Physics

## Excess electronic recoil events in XENON1T

E. Aprile,<sup>1</sup> J. Aalbers,<sup>2</sup> F. Agostini,<sup>3</sup> M. Alfonsi,<sup>4</sup> L. Althueser,<sup>5</sup> F. D. Amaro,<sup>6</sup> V. C. Antochi,<sup>2</sup> E. Angelino,<sup>7</sup> J. R. Angevaare,<sup>8</sup> F. Arneodo,<sup>9</sup> D. Barge,<sup>2</sup> L. Baudis,<sup>10</sup> B. Bauermeister,<sup>2</sup> L. Bellagamba,<sup>3</sup> M. L. Benabderahmane,<sup>9</sup> T. Berger,<sup>11</sup> A. Brown,<sup>10</sup> E. Brown,<sup>11</sup> S. Bruennner,<sup>8</sup> G. Bruno,<sup>9</sup> R. Budnik,<sup>12,\*</sup> C. Capelli,<sup>10</sup> J. M. R. Cardoso,<sup>6</sup> D. Cichon,<sup>13</sup> B. Cimmino,<sup>14</sup> M. Clark,<sup>15</sup> D. Coderre,<sup>16</sup> A. P. Colijn,<sup>8,†</sup> J. Conrad,<sup>2</sup> J. P. Cussonneau,<sup>17</sup> M. P. Decowski,<sup>8</sup> A. Depoian,<sup>15</sup> P. Di Gangi,<sup>3</sup> A. Di Giovanni,<sup>9</sup> R. Di Stefano,<sup>14</sup> S. Diglio,<sup>17</sup> A. Elykov,<sup>16</sup> G. Eurin,<sup>13</sup> A. D. Ferella,<sup>18,19</sup> W. Fulgione,<sup>7,19</sup> P. Gaemers,<sup>8</sup> R. Gaior,<sup>20</sup> M. Galloway,<sup>10,‡</sup> F. Gao,<sup>1</sup> L. Grandi,<sup>21</sup> C. Hasterok,<sup>13</sup> C. Hils,<sup>4</sup> K. Hiraide,<sup>22</sup> L. Hoetzsch,<sup>13</sup> J. Howlett,<sup>1</sup> M. Iacovacci,<sup>14</sup> Y. Itow,<sup>23</sup> F. Joerg,<sup>13</sup> N. Kato,<sup>22</sup> S. Kazama,<sup>23,§</sup> M. Kobayashi,<sup>1</sup> G. Koltman,<sup>12</sup> A. Kopec,<sup>15</sup> H. Landsman,<sup>12</sup> R. F. Lang,<sup>15</sup> L. Levinson,<sup>12</sup> Q. Lin,<sup>1</sup> S. Lindemann,<sup>16</sup> M. Lindner,<sup>13</sup> F. Lombardi,<sup>6</sup> J. Long,<sup>21</sup> J. A. M. Lopes,<sup>6,||</sup> E. López Fune,<sup>20</sup> C. Macolino,<sup>24</sup> J. Mahlstedt,<sup>2</sup> A. Mancuso,<sup>3</sup> L. Manenti,<sup>9</sup> A. Manfredini,<sup>10</sup> F. Marignetti,<sup>14</sup> T. Marrodán Undagoitia,<sup>13</sup> K. Martens,<sup>22</sup> J. Masbou,<sup>17</sup> D. Masson,<sup>16</sup> S. Mastroianni,<sup>14</sup> M. Messina,<sup>19</sup> K. Miuchi,<sup>25</sup> K. Mizukoshi,<sup>25</sup> A. Molinario,<sup>19</sup> K. Morå,<sup>1,2</sup> S. Moriyama,<sup>22</sup> Y. Mosbacher,<sup>12</sup> M. Murra,<sup>5</sup> J. Naganoma,<sup>19</sup> K. Ni,<sup>26</sup> U. Oberlack,<sup>4</sup> K. Odgers,<sup>11</sup> J. Palacio,<sup>13,17</sup> B. Pelssers,<sup>2</sup> R. Peres,<sup>10</sup> J. Pienaar,<sup>21</sup> V. Pizzella,<sup>13</sup> G. Plante,<sup>1</sup> J. Qin,<sup>15</sup> H. Qiu,<sup>12</sup> D. Ramírez García,<sup>16</sup> S. Reichard,<sup>10</sup> A. Rocchetti,<sup>16</sup> N. Rupp,<sup>13</sup> J. M. F. dos Santos,<sup>6</sup> G. Sartorelli,<sup>3</sup> N. Šarčević,<sup>16</sup> M. Scheibelhut,<sup>4</sup> J. Schreiner,<sup>13</sup> D. Schulte,<sup>5</sup> M. Schumann,<sup>16</sup> L. Scotto Lavina,<sup>20</sup> M. Selvi,<sup>3</sup> F. Semeria,<sup>3</sup> P. Shagin,<sup>27</sup> E. Shockley,<sup>21,¶</sup> M. Silva,<sup>6</sup> H. Simgen,<sup>13</sup> A. Takeda,<sup>22</sup> C. Therreau,<sup>17</sup> D. Thers,<sup>17</sup> F. Toschi,<sup>16</sup> G. Trinchero,<sup>7</sup> C. Tunnell,<sup>27</sup> M. Vargas,<sup>5</sup> G. Volta,<sup>10</sup> H. Wang,<sup>28</sup> Y. Wei,<sup>26</sup> C. Weinheimer,<sup>5</sup> M. Weiss,<sup>12</sup> D. Wenz,<sup>4</sup> C. Wittweg,<sup>5</sup> Z. Xu,<sup>1</sup> M. Yamashita,<sup>23,22</sup> J. Ye,<sup>26,\*\*</sup> G. Zavattini,<sup>3,††</sup> Y. Zhang,<sup>1</sup> T. Zhu,<sup>1</sup> and J. P. Zopounidis,<sup>20</sup>

(XENON Collaboration)<sup>††</sup>

X. Mougeot<sup>29</sup>

# An excess of events

PRD 102 (2020) 072004

PHYSICAL REVIEW D 102, 072004 (2020)


Featured in Physics

## Excess electronic recoil events in XENON1T

E. Aprile,<sup>1</sup> J. Aalbers,<sup>2</sup> F. Agostini,<sup>3</sup> M. Alfonsi,<sup>4</sup> L. Althueser,<sup>5</sup> F. D. Amaro,<sup>6</sup> V. C. Antochi,  
J. R. Angevaare,<sup>8</sup> F. Arneodo,<sup>9</sup> D. Barge,<sup>2</sup> L. Baudis,<sup>10</sup> B. Bauermeister,<sup>2</sup> L. Bellagamba,<sup>3</sup> M. L.  
T. Berger,<sup>11</sup> A. Brown,<sup>10</sup> E. Brown,<sup>11</sup> S. Bruenner,<sup>8</sup> G. Bruno,<sup>9</sup> R. Budnik,<sup>12,\*</sup> C. Capelli,<sup>10</sup> J. M. R. C.  
B. Cimmino,<sup>14</sup> M. Clark,<sup>15</sup> D. Coderre,<sup>16</sup> A. P. Colijn,<sup>8,†</sup> J. Conrad,<sup>2</sup> J. P. Cussonneau,<sup>17</sup> M. P. Deco  
P. Di Gangi,<sup>3</sup> A. Di Giovanni,<sup>9</sup> R. Di Stefano,<sup>14</sup> S. Diglio,<sup>17</sup> A. Elykov,<sup>16</sup> G. Eurin,<sup>13</sup> A. D. Ferella,  
P. Gaemers,<sup>8</sup> R. Gaior,<sup>20</sup> M. Galloway<sup>10,‡</sup>, F. Gao,<sup>1</sup> L. Grandi,<sup>21</sup> C. Hasterok,<sup>13</sup> C. Hils,<sup>4</sup> K. Hira  
J. Howlett,<sup>1</sup> M. Iacovacci,<sup>14</sup> Y. Itow,<sup>23</sup> F. Joerg,<sup>13</sup> N. Kato,<sup>22</sup> S. Kazama,<sup>23,§</sup> M. Kobayashi,<sup>1</sup> G. Ko  
H. Landsman,<sup>12</sup> R. F. Lang,<sup>15</sup> L. Levinson,<sup>12</sup> Q. Lin,<sup>1</sup> S. Lindemann,<sup>16</sup> M. Lindner,<sup>13</sup> F. Lomb  
J. A. M. Lopes,<sup>6,||</sup> E. López Fune,<sup>20</sup> C. Macolino,<sup>24</sup> J. Mahlstedt,<sup>2</sup> A. Mancuso,<sup>3</sup> L. Manenti,<sup>9</sup> A. Manfredini,<sup>10</sup>  
F. Marignetti,<sup>14</sup> T. Marrodán Undagoitia,<sup>13</sup> K. Martens,<sup>22</sup> J. Masbou,<sup>17</sup> D. Masson,<sup>16</sup> S. Mastroianni,<sup>14</sup> M. Messina,<sup>19</sup>  
K. Miuchi,<sup>25</sup> K. Mizukoshi,<sup>25</sup> A. Molinario,<sup>19</sup> K. Morå,<sup>1,2</sup> S. Moriyama,<sup>22</sup> Y. Mosbacher,<sup>12</sup> M. Murra,<sup>5</sup> J. Naganoma,<sup>19</sup>  
K. Ni,<sup>26</sup> U. Oberlack,<sup>4</sup> K. Odgers,<sup>11</sup> J. Palacio,<sup>13,17</sup> B. Pelssers,<sup>2</sup> R. Peres,<sup>10</sup> J. Pienaar,<sup>21</sup> V. Pizzella,<sup>13</sup> G. Plante,<sup>1</sup> J. Qin,<sup>15</sup>  
H. Qiu,<sup>12</sup> D. Ramírez García,<sup>16</sup> S. Reichard,<sup>10</sup> A. Rocchetti,<sup>16</sup> N. Rupp,<sup>13</sup> J. M. F. dos Santos,<sup>6</sup> G. Sartorelli,<sup>3</sup> N. Šarčević,<sup>16</sup>  
M. Scheibelhut,<sup>4</sup> J. Schreiner,<sup>13</sup> D. Schulte,<sup>5</sup> M. Schumann,<sup>16</sup> L. Scotto Lavina,<sup>20</sup> M. Selvi,<sup>3</sup> F. Semeria,<sup>3</sup> P. Shagin,<sup>27</sup>  
E. Shockley<sup>10,¶</sup>, M. Silva,<sup>6</sup> H. Simgen,<sup>13</sup> A. Takeda,<sup>22</sup> C. Therreau,<sup>17</sup> D. Thers,<sup>17</sup> F. Toschi,<sup>16</sup> G. Trinchero,<sup>7</sup> C. Tunnell,<sup>27</sup>  
M. Vargas,<sup>5</sup> G. Volta,<sup>10</sup> H. Wang,<sup>28</sup> Y. Wei,<sup>26</sup> C. Weinheimer,<sup>5</sup> M. Weiss,<sup>12</sup> D. Wenz,<sup>4</sup> C. Wittweg,<sup>5</sup> Z. Xu,<sup>1</sup>  
M. Yamashita,<sup>23,22</sup> J. Ye<sup>26,\*\*</sup>, G. Zavattini,<sup>3,††</sup> Y. Zhang,<sup>1</sup> T. Zhu,<sup>1</sup> and J. P. Zopounidis,<sup>20</sup>

(XENON Collaboration)<sup>††</sup>

X. Mougeot<sup>29</sup>



Released on  
**17 June**

# An excess of events

PRD 102 (2020) 072004

PHYSICAL REVIEW D 102, 072004 (2020)


Featured in Physics

## Excess electronic recoil events in XENON1T

E. Aprile,<sup>1</sup> J. Aalbers,<sup>2</sup> F. Agostini,<sup>3</sup> M. Alfonsi,<sup>4</sup> L. Althueser,<sup>5</sup> F. D. Amaro,<sup>6</sup> V. C. Antochi,  
J. R. Angevaare,<sup>8</sup> F. Arneodo,<sup>9</sup> D. Barge,<sup>2</sup> L. Baudis,<sup>10</sup> B. Bauermeister,<sup>2</sup> L. Bellagamba,<sup>3</sup> M. L.  
T. Berger,<sup>11</sup> A. Brown,<sup>10</sup> E. Brown,<sup>11</sup> S. Bruenner,<sup>8</sup> G. Bruno,<sup>9</sup> R. Budnik,<sup>12,\*</sup> C. Capelli,<sup>10</sup> J. M. R. C.  
B. Cimmino,<sup>14</sup> M. Clark,<sup>15</sup> D. Coderre,<sup>16</sup> A. P. Colijn,<sup>8,†</sup> J. Conrad,<sup>2</sup> J. P. Cussonneau,<sup>17</sup> M. P. Deco  
P. Di Gangi,<sup>3</sup> A. Di Giovanni,<sup>9</sup> R. Di Stefano,<sup>14</sup> S. Diglio,<sup>17</sup> A. Elykov,<sup>16</sup> G. Eurin,<sup>13</sup> A. D. Ferella,  
P. Gaemers,<sup>8</sup> R. Gaior,<sup>20</sup> M. Galloway<sup>10,‡</sup>, F. Gao,<sup>1</sup> L. Grandi,<sup>21</sup> C. Hasterok,<sup>13</sup> C. Hils,<sup>4</sup> K. Hira  
J. Howlett,<sup>1</sup> M. Iacovacci,<sup>14</sup> Y. Itow,<sup>23</sup> F. Joerg,<sup>13</sup> N. Kato,<sup>22</sup> S. Kazama,<sup>23,§</sup> M. Kobayashi,<sup>1</sup> G. Ko  
H. Landsman,<sup>12</sup> R. F. Lang,<sup>15</sup> L. Levinson,<sup>12</sup> Q. Lin,<sup>1</sup> S. Lindemann,<sup>16</sup> M. Lindner,<sup>13</sup> F. Lomb  
J. A. M. Lopes,<sup>6,||</sup> E. López Fune,<sup>20</sup> C. Macolino,<sup>24</sup> J. Mahlstedt,<sup>2</sup> A. Mancuso,<sup>3</sup> L. Manenti,<sup>9</sup> A. Manfredini,<sup>10</sup>  
F. Marignetti,<sup>14</sup> T. Marrodán Undagoitia,<sup>13</sup> K. Martens,<sup>22</sup> J. Masbou,<sup>17</sup> D. Masson,<sup>16</sup> S. Mastroianni,<sup>14</sup> M. Messina,<sup>19</sup>  
K. Miuchi,<sup>25</sup> K. Mizukoshi,<sup>25</sup> A. Molinario,<sup>19</sup> K. Morå,<sup>1,2</sup> S. Moriyama,<sup>22</sup> Y. Mosbacher,<sup>12</sup> M. Murra,<sup>5</sup> J. Naganoma,<sup>19</sup>  
K. Ni,<sup>26</sup> U. Oberlack,<sup>4</sup> K. Odgers,<sup>11</sup> J. Palacio,<sup>13,17</sup> B. Pelssers,<sup>2</sup> R. Peres,<sup>10</sup> J. Pienaar,<sup>21</sup> V. Pizzella,<sup>13</sup> G. Plante,<sup>1</sup> J. Qin,<sup>15</sup>  
H. Qiu,<sup>12</sup> D. Ramírez García,<sup>16</sup> S. Reichard,<sup>10</sup> A. Rocchetti,<sup>16</sup> N. Rupp,<sup>13</sup> J. M. F. dos Santos,<sup>6</sup> G. Sartorelli,<sup>3</sup> N. Šarčević,<sup>16</sup>  
M. Scheibelhut,<sup>4</sup> J. Schreiner,<sup>13</sup> D. Schulte,<sup>5</sup> M. Schumann,<sup>16</sup> L. Scotto Lavina,<sup>20</sup> M. Selvi,<sup>3</sup> F. Semeria,<sup>3</sup> P. Shagin,<sup>27</sup>  
E. Shockley<sup>10,¶</sup>, M. Silva,<sup>6</sup> H. Simgen,<sup>13</sup> A. Takeda,<sup>22</sup> C. Therreau,<sup>17</sup> D. Thers,<sup>17</sup> F. Toschi,<sup>16</sup> G. Trinchero,<sup>7</sup> C. Tunnell,<sup>27</sup>  
M. Vargas,<sup>5</sup> G. Volta,<sup>10</sup> H. Wang,<sup>28</sup> Y. Wei,<sup>26</sup> C. Weinheimer,<sup>5</sup> M. Weiss,<sup>12</sup> D. Wenz,<sup>4</sup> C. Wittweg,<sup>5</sup> Z. Xu,<sup>1</sup>  
M. Yamashita,<sup>23,22</sup> J. Ye<sup>26,\*\*</sup>, G. Zavattini,<sup>3,††</sup> Y. Zhang,<sup>1</sup> T. Zhu,<sup>1</sup> and J. P. Zopounidis,<sup>20</sup>

(XENON Collaboration)<sup>††</sup>

X. Mougeot<sup>29</sup>



## Search for New Physics with Electronic-Recoil Events in XENON1T

Evan Shockley

The University of Chicago & KICP

For the XENON collaboration + X. Mougeot

LNGS Webinar


June 17th, 2020

Released on  
**17 June**

Cited  
**157 times**


EDITORS' PICK | 5326 views | Jun 17, 2020, 10:00am EDT

# Is It Dark Matter? Mysterious Signal Goes 'Bump' In World's Most Sensitive Detector

**Ethan Siegel** Senior Contributor  
**Starts With A Bang** Contributor Group 

Science

The Universe is out there, waiting for you to discover it.



**Il Messaggero**

Ancora una volta la parola che più si accosta al termine Fisica è Italia. E seppur con la dovuta cautela gli scienziati vanno a passo di piombo, gli avvenimenti accaduti nei **Laboratori del Gran Sasso** prefigurano scene

# An excess of press articles



Sign in

Home

News

Sport

Reel

Worklife

T

NEWS

Home | US Election | Coronavirus | Video | World | UK | Business | Tech | Science | Stories | Entertainment

Science

## Dark matter hunt yields unexplained signal

By Paul Rincon  
 Science editor, BBC News website

Le Monde

de  
Consulter le journal

ACTUALITÉS

ÉCONOMIE

VIDÉOS

OPINIONS

CULTURE

M LE MAG

SCIENCES • PHYSIQUE

Favoris Pa

## Physique des particules : des détections déroutantes en Italie

L'expérience Xenon1T, installée dans le laboratoire souterrain de Gran Sasso pour traquer la matière noire a enregistré un signal inattendu. S'agit-il d'une particule encore jamais observée ou d'une contamination du dispositif ?

Par Nathaniel Herzberg · Publié le 19 juin 2020 à 14h23 - Mis à jour le 22 juin 2020 à 15h59

Quanta magazine

Physics Mathematics Biology Computer Science All Articles

ABSTRACTIONS BLOG

## Dark Matter Experiment Finds Unexplained Signal

Researchers say there are three possible explanations for the anomalous data. One is mundane. Two would revolutionize physics.

11

1

GLI ESPERIMENTI SOTTO IL GRAN SASSO


## Laboratorio del Gran Sasso, registrati segnali anomali nella caccia alla materia oscura: sono gli assioni?

The New York Times

OUT THERE

## Seeking Dark Matter, They Detected Another Mystery

Do signals from beneath an Italian mountain herald a revolution in physics?



JUNE 17, 2020

## Observation of excess events in the XENON1T dark matter experiment

by Kavli Institute for the Physics and Mathematics of the Universe

Science

Contents News Careers Journals

Read our COVID-19 research and news.

SHARE

f t in e 3

  
The XENON1T detector, which ran from 2016 to 2018, may have seen signs of exotic particles—or not. ENRICO SACCHETTI/SCIENCE SOURCE

Dark matter hunters' inconclusive signal grabs headlines

By Adrian Cho | Jun. 17, 2020, 5:55 PM

SciTechDaily

BIOLOGY CHEMISTRY EARTH HEALTH PHYSICS SCIENCE SPACE TECHNOLOGY

HOT TOPICS OCTOBER 21, 2020 | RESEARCHERS USE "ARTIFICIAL GRAMMARS" TO SHOW BUILDING BLOCKS OF LANGUAGE EVOLVED 30-40 MILLION YEARS AGO

HOME PHYSICS NEWS

Exotic Dark Matter Detector Deep in an Italian Mountainside Picks Up Unexplained New Signal

TOPICS: Dark Matter Particle Physics Popular University Of Chicago

By MAGGIE HUDSON, UNIVERSITY OF CHICAGO JUNE 18, 2020

MENU CERCA




## Materia Oscura, mistero sui nuovi eventi scoperti nei laboratori del Gran Sasso

SCIENZA

Giovedì 18 Giugno 2020 di Enzo Vitale


# The observation

PRD 102 (2020) 072004




# The observation

PRD 102 (2020) 072004




# The observation

PRD 102 (2020) 072004




# The observation

PRD 102 (2020) 072004



# The observation


PRD 102 (2020) 072004



- ▶ Carefully selected **electronic recoil data** from XENON1T Science Run 1
- ▶ Comprehensive **background model**
- ▶ Observed **285 events**
- ▶ Expected  $(232 \pm 15)$  events in 1–7 keV  
→  $\sim 3.3\sigma$  Poissonian fluctuation!
- ▶ What is going on here?

# Data selection


PRD 100 (2019) 052014  
PRD 102 (2020) 072004



- ▶ Science Run 1 (SR1, Feb. 2017– Feb. 2018) data → **226.9 d effective live time**  
(+ 24.4 d cross-check data from post-SR1 SR2 with 20% less ER background)

# Data selection

PRD 100 (2019) 052014  
PRD 102 (2020) 072004





- ▶ Science Run 1 (SR1, Feb. 2017– Feb. 2018) data → **226.9 d effective live time**  
(+ 24.4 d cross-check data from post-SR1 SR2 with 20% less ER background)
- ▶ Surface and material background mitigation → **1 042 kg LXe fiducial volume**

**0.65 t × yr  
total  
exposure**

# Data selection

PRD 100 (2019) 052014  
PRD 102 (2020) 072004




- ▶ Science Run 1 (SR1, Feb. 2017– Feb. 2018) data → **226.9 d effective live time**  
(+ 24.4 d cross-check data from post-SR1 SR2 with 20% less ER background)
- ▶ Surface and material background mitigation → **1 042 kg LXe fiducial volume**
- ▶ Selection of high S2 threshold, single scatter events (+ other requirements) **in 1–210 keV**

**0.65 t × yr  
total  
exposure**

# Background model $B_0$


PRD 102 (2020) 072004




- **10-component** background model (dominated by  $^{214}\text{Pb}$   $\beta$  decay from  $^{222}\text{Rn}$  emanation)

# Background model $B_0$

PRD 102 (2020) 072004




## Intrinsic sources




- **10-component** background model (dominated by  $^{214}\text{Pb}$   $\beta$  decay from  $^{222}\text{Rn}$  emanation)

# Background model $B_0$

PRD 102 (2020) 072004




## Intrinsic sources



- **10-component** background model (dominated by  $^{214}\text{Pb}$   $\beta$  decay from  $^{222}\text{Rn}$  emanation)

# Background model $B_0$


PRD 102 (2020) 072004



## Intrinsic sources




## Neutron-activated



- **10-component** background model (dominated by  $^{214}\text{Pb}$   $\beta$  decay from  $^{222}\text{Rn}$  emanation)

# Background model $B_0$

PRD 102 (2020) 072004



## Intrinsic sources




## Neutron-activated



- **10-component** background model (dominated by  $^{214}\text{Pb}$   $\beta$  decay from  $^{222}\text{Rn}$  emanation)

# Background model $B_0$

PRD 102 (2020) 072004



## Intrinsic sources



## Neutron-activated



## Detector materials $\gamma$


## Solar neutrinos

*both continuous spectra*

- ▶ **10-component** background model (dominated by  $^{214}\text{Pb}$   $\beta$  decay from  $^{222}\text{Rn}$  emanation)
- ▶ Theoretical and GEANT4 spectra, **convolved with detector effects** (resolution, efficiency)

# Data-model matching


PRD 102 (2020) 072004



- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)

# Data-model matching


PRD 102 (2020) 072004



- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)
- ▶ Very good fit over the studied energy range (1–210 keV), **but look closer...**

# Data-model matching


PRD 102 (2020) 072004



- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)
- ▶ Very good fit over the studied energy range (1–210 keV), **but look closer...**

# Data-model matching

PRD 102 (2020) 072004




- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)
- ▶ Very good fit over the studied energy range (1–210 keV), **but look closer...**

Lowest background rate ever  
achieved in 1–30 keV!  
**( $76 \pm 2$ ) events/( $t \times yr \times \text{keV}$ )**

# Data-model matching

PRD 102 (2020) 072004




- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)
- ▶ Very good fit over the studied energy range (1–210 keV), **but look closer...**

Lowest background rate ever achieved in 1–30 keV!  
 $(76 \pm 2)$  events/( $t \times yr \times keV$ )



# Data-model matching

PRD 102 (2020) 072004



- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)
- ▶ Very good fit over the studied energy range (1–210 keV), **but look closer...**

Lowest background rate ever achieved in 1–30 keV!  
 $(76 \pm 2)$  events/ $(t \times yr \times \text{keV})$




Statistical  
fluke?



Systematic  
effects?

# Data-model matching

PRD 102 (2020) 072004



- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)
- ▶ Very good fit over the studied energy range (1–210 keV), **but look closer...**


Lowest background rate ever achieved in 1–30 keV!  
 $(76 \pm 2)$  events/ $(t \times yr \times \text{keV})$

**Statistical fluke?**      **Systematic effects?**

**Unexplored background?**

# Data-model matching

PRD 102 (2020) 072004




- ▶ Fitting with an **unbinned profile likelihood** (data binned for display)
- ▶ Very good fit over the studied energy range (1–210 keV), **but look closer...**

Lowest background rate ever achieved in 1–30 keV!  
 $(76 \pm 2)$  events/ $(t \times yr \times \text{keV})$



# Statistical fluke?


PRD 102 (2020) 072004



# Statistical fluke?




PRD 102 (2020) 072004



# Statistical fluke?



PRD 102 (2020) 072004





- KDE with energy resolution as width → **low-energy excess still significant**, the rest no

# Statistical fluke?



PRD 102 (2020) 072004





- ▶ KDE with energy resolution as width → **low-energy excess still significant**, the rest no
- ▶ Display binning change → bin width smaller than energy resolution, **low-energy excess robustly visible across several bins**, the rest no

# Statistical fluke?



PRD 102 (2020) 072004




UNLIKELY

- ▶ KDE with energy bins → **low-energy excess still significant**, the rest no
- ▶ Display binning → bin width smaller than energy resolution, **low-energy excess robustly visible across several bins**, the rest no

# Systematic effects?




PRD 102 (2020) 072004



# Systematic effects?




PRD 102 (2020) 072004



# Systematic effects?

PRD 102 (2020) 072004


- ▶ Excess close to but **not at the analysis threshold**  
( $\sim 85\%$  detection efficiency at 2–3 keV)
- ▶ Still present if threshold is **doubled**



# Systematic effects?

PRD 102 (2020) 072004



- ▶ Excess close to but **not at the analysis threshold**  
( $\sim 85\%$  detection efficiency at 2–3 keV)
- ▶ Still present if threshold is **doubled**
- ▶ Efficiency verified with high-statistic  $^{220}\text{Rn}$  calibration  
data → **background model validated**



# Systematic effects?

PRD 102 (2020) 072004

- ▶ Excess close to but **not at the analysis threshold**  
( $\sim 85\%$  detection efficiency at 2–3 keV)
- ▶ Still present if threshold is **doubled**
- ▶ Efficiency verified with high-statistic  $^{220}\text{Rn}$  calibration data → **background model validated**




- ▶ Surface events → **absent from the search region** thanks to fiducialisation
- ▶ Accidental Coincidences (AC events) → **tightly constrained** and well-understood (S1, S2) signature

# Systematic effects?




PRD 102 (2020) 072004


- ▶ Excess close to but **not at the analysis threshold**  
( $\sim 85\%$  detection efficiency at 2–3 keV)
- ▶ Still present if threshold is **doubled**
- ▶ Efficiency verified with high-statistic  
data → **background model**




- Rare events → **absent from the search region**  
thanks to fiducialisation
- ▶ Accidental Coincidences (AC events) → **tightly constrained** and well-understood (S1, S2) signature




# Unexplored background?




PRD 102 (2020) 072004




# Unexplored background?




PRD 102 (2020) 072004



# Many background sources




# Many background sources




- ▶ Through  $> 1$  km of rock
- ▶ **Muons** → too high energy
- ▶ **Neutrinos & dark particles**

# Many background sources



- ▶ Through  $> 1$  km of rock  
**Muons** → too high energy  
**Neutrinos & dark particles**
- ▶ Through cms of metal/LXe  
**Neutrons** → NR, not ER  
 $\beta$  &  $\gamma$  → too high energy


# Many background sources



# The tritium hypothesis

NDS 130 (2015) 1–20  
PRD 102 (2020) 072004



- ▶ **Never observed before** as background in LXe TPCs
- ▶  $\beta^-$  emitter  $\rightarrow Q_\beta = 18.6 \text{ keV}$ ,  $t_{1/2} = 12.3 \text{ yr}$
- ▶ Cosmogenic activation? Atmospheric abundance?



# The tritium hypothesis

NDS 130 (2015) 1–20  
PRD 102 (2020) 072004

- Never observed before as background in LXe TPCs
- $\beta^-$  emitter  $\rightarrow Q_\beta = 18.6 \text{ keV}$ ,  $t_{1/2} = 12.3 \text{ yr}$
- Cosmogenic activation? Atmospheric abundance?




Tritium hypothesis  
favoured over  $B_0$  at  **$3.2\sigma$**

- Fitted rate  $\rightarrow (159 \pm 51) \text{ events}/(t \times \text{yr})$
- ${}^3\text{H}/\text{Xe} = (6.2 \pm 2.0) \times 10^{-25} \text{ mol/mol}$
- $\rightarrow \lesssim 3 \text{ tritium atoms/kg of xenon}$

# Tritium from the cosmos?

PRD 102 (2020) 072004


1. Xenon gas storage above ground → ~ 32 tritium atoms/(kg × d) → **tritiated water (HTO)** with ~ 1 ppm water impurities



# Tritium from the cosmos?

PRD 102 (2020) 072004


1. Xenon gas storage above ground →  $\sim 32$  tritium atoms/(kg × d) → **tritiated water (HTO)** with  $\sim 1$  ppm water impurities
2. Xenon gas moved underground → **tritium decay**, no more cosmogenic activation



# Tritium from the cosmos?

PRD 102 (2020) 072004


1. Xenon gas storage above ground →  $\sim 32$  tritium atoms/(kg × d) → **tritiated water (HTO)** with  $\sim 1$  ppm water impurities
2. Xenon gas moved underground → **tritium decay**, no more cosmogenic activation
3. Xenon filled into its cold storage vessel → water condensation (including HTO) and “capture” by the vessel walls  
→  **$\times 4000$  HTO reduction**



# Tritium from the cosmos?

PRD 102 (2020) 072004


1. Xenon gas storage above ground →  $\sim 32$  tritium atoms/(kg × d) → **tritiated water (HTO)** with  $\sim 1$  ppm water impurities
2. Xenon gas moved underground → **tritium decay**, no more cosmogenic activation
3. Xenon filled into its cold storage vessel → water condensation (including HTO) and “capture” by the vessel walls  
→  **$\times 4000$  HTO reduction**
4. Further tritium decay



# Tritium from the cosmos?

PRD 102 (2020) 072004

1. Xenon gas storage above ground →  $\sim 32$  tritium atoms/(kg × d) → **tritiated water (HTO)** with  $\sim 1$  ppm water impurities
2. Xenon gas moved underground → **tritium decay**, no more cosmogenic activation
3. Xenon filled into its cold storage vessel → water condensation (including HTO) and “capture” by the vessel walls  
→ **× 4000 HTO reduction**
4. Further tritium decay
5. Purification and detector filling  
→ **99.99% water removal**  
(including HTO)



# Tritium from the cosmos?

PRD 102 (2020) 072004

1. Xenon gas storage above ground → ~ 32 tritium atoms  
with ~ 1 ppm water impurities
  2. Xenon gas moved underground → tritium and water ( $\text{HTO}$ )  
decay
  3. Xenon filled into its cold storage vessel → water condensation  
(including  $\text{HTO}$ ) and “tritium rain” on the vessel walls  
→  $\times 4000$  times more tritium
  4. Further tritium decay
  5. Purification and separation  
→ **99.99% water**  
(including  $\text{HTO}$ )
- UNLIKELY**
-

# Tritium from the atmosphere?

PRD 102 (2020) 072004


- ▶ Atmospheric abundance → tritium stored as **tritiated water (HTO) or hydrogen (HT)** in detector materials, then **emanating in xenon**
- ▶  $\text{HTO}/\text{H}_2\text{O} = (5\text{--}10) \times 10^{-18}$  mol/mol, **assuming** the same value for  $\text{HT}/\text{H}_2$   
→  **$(\text{H}_2\text{O} + \text{H}_2)/\text{Xe} \gtrsim 30 \text{ ppb}$**  required to make up the tritium-fitted excess

# Tritium from the atmosphere?

PRD 102 (2020) 072004

- ▶ Atmospheric abundance → tritium stored as **tritiated water (HTO)** or **hydrogen (HT)** in detector materials, then **emanating in xenon**
- ▶  $\text{HTO}/\text{H}_2\text{O} = (5\text{--}10) \times 10^{-18}$  mol/mol, **assuming** the same value for  $\text{HT}/\text{H}_2$   
→  **$(\text{H}_2\text{O} + \text{H}_2)/\text{Xe} \gtrsim 30$  ppb** required to make up the tritium-fitted excess


## Tritiated water (HTO)



# Tritium from the atmosphere?

PRD 102 (2020) 072004



- ▶ Atmospheric abundance → tritium stored as **tritiated water (HTO)** or **hydrogen (HT)** in detector materials, then **emanating in xenon**
- ▶  $\text{HTO}/\text{H}_2\text{O} = (5\text{--}10) \times 10^{-18}$  mol/mol, **assuming** the same value for  $\text{HT}/\text{H}_2$   
→  **$(\text{H}_2\text{O} + \text{H}_2)/\text{Xe} \gtrsim 30$  ppb** required to make up the tritium-fitted excess



# Tritium from the atmosphere?

PRD 102 (2020) 072004



- Atmospheric abundance → tritium stored as **tritiated water (HTO)** or **hydrogen (HT)** in detector materials, then **emanating in xenon**
- $\text{HTO}/\text{H}_2\text{O} = (5\text{--}10) \times 10^{-18}$  mol/mol, **assuming** the same value for  $\text{HT}/\text{H}_2$   
→  $(\text{H}_2\text{O} + \text{H}_2)/\text{Xe} \gtrsim 30$  ppb required to make up the tritium-fitted excess



# Tritium from the atmosphere?

PRD 102 (2020) 072004


- Atmospheric abundance → tritium stored as **tritiated water (HTO)** or **hydrogen (HT)** in detector materials, then **emanating in xenon**
- $\text{HTO}/\text{H}_2\text{O} = (5\text{--}10) \times 10^{-18}$  mol/mol, **assuming** the same value for  $\text{HT}/\text{H}_2$   
→  $(\text{H}_2\text{O} + \text{H}_2)/\text{Xe} \gtrsim 30$  ppb required to make up the tritium-fitted excess



# New physics?



PRD 102 (2020) 072004




# New physics?



PRD 102 (2020) 072004



Statistical  
fluke?




# New physics?



PRD 102 (2020) 072004

UNLIKELY  
real  
noise?




# New physics?



PRD 102 (2020) 072004

UNLIKELY  
real  
noise?




Systematic  
effects?

# New physics?



PRD 102 (2020) 072004

UNLIKELY  
Irreducible background?



UNLIKELY  
Systematic effects?


# New physics?



PRD 102 (2020) 072004

UNLIKELY  
signal?  
Fake?

Unexplored  
background?




UNLIKELY  
signals?  
Effects?

# New physics?



PRD 102 (2020) 072004

UNLIKELY  
Is it real?  
Is it fake?  
Is it real?  
Is it fake?  
Is it real?  
Is it fake?  
Is it real?  
Is it fake?




UNLIKELY  
Is it systematic?  
Is it statistical?  
Is it systematic?  
Is it statistical?

# New physics?



PRD 102 (2020) 072004

UNLIKELY  
Is it real?  
Is it fake?  
Is it signal?  
Is it background?



UNLIKELY  
Is it systematic?  
Is it statistical?  
Is it effects?

New  
physics?  
Many models,  
three approaches  
considered  
in our paper

# The axion hypothesis

PRD 102 (2020) 072004


- ▶ Hypothetical particle suggested as a **QCD strong  $CP$  problem solution**
- ▶ Motivated **Dark Matter candidates** ( $\ll \text{keV}$ ) and possibly **produced in the Sun** ( $\sim \text{keV}$ )

# The axion hypothesis

PRD 102 (2020) 072004

- ▶ Hypothetical particle suggested as a **QCD strong CP problem solution**
- ▶ Motivated **Dark Matter candidates** ( $\ll \text{keV}$ ) and possibly **produced in the Sun** ( $\sim \text{keV}$ )

Atomic recombination and de-excitation,  
Bremsstrahlung and Compton (ABC)  
 $\propto$  Axion-electron coupling ( $g_{ae}$ )<sup>2</sup>




# The axion hypothesis

PRD 102 (2020) 072004

- ▶ Hypothetical particle suggested as a **QCD strong CP problem solution**
- ▶ Motivated **Dark Matter candidates** ( $\ll \text{keV}$ ) and possibly **produced in the Sun** ( $\sim \text{keV}$ )

Atomic recombination and de-excitation,  
Bremsstrahlung and Compton (ABC)  
 $\propto$  Axion-electron coupling ( $g_{ae}$ )<sup>2</sup>


Primakoff conversion of  
photons to axions  
 $\propto$  Axion-photon coupling ( $g_{a\gamma}$ )<sup>2</sup>



# The axion hypothesis

PRD 102 (2020) 072004


- ▶ Hypothetical particle suggested as a **QCD strong CP problem solution**
- ▶ Motivated **Dark Matter candidates** ( $\ll \text{keV}$ ) and possibly **produced in the Sun** ( $\sim \text{keV}$ )



# The axion hypothesis

PRD 102 (2020) 072004

- ▶ Hypothetical particle suggested as a **QCD strong CP problem solution**
- ▶ Motivated **Dark Matter candidates** ( $\ll \text{keV}$ ) and possibly **produced in the Sun** ( $\sim \text{keV}$ )




Detection via  
axioelectric effect  
 $\propto (g_{ae})^2$   
⊗  
Reconstruction  
effects


# The axion hypothesis

PRD 102 (2020) 072004

- ▶ Hypothetical particle suggested as a **QCD strong CP problem solution**
- ▶ Motivated **Dark Matter candidates** ( $\ll \text{keV}$ ) and possibly **produced in the Sun** ( $\sim \text{keV}$ )




Detection via  
axioelectric effect  
 $\propto (g_{ae})^2$   
 $\otimes$   
Reconstruction  
effects



# Solar axion results


PRD 102 (2020) 072004



► Simultaneous search for **ABC**,  
**Primakoff** and  **$^{57}\text{Fe}$**  axions  
(unconstrained components in the fit)

# Solar axion results

PRD 102 (2020) 072004




- ▶ Simultaneous search for **ABC**, **Primakoff** and  **$^{57}\text{Fe}$**  axions  
(unconstrained components in the fit)


Axion hypothesis  
favoured over  $B_0$  at  **$3.4\sigma$**

# Solar axion results

PRD 102 (2020) 072004




- ▶ Simultaneous search for **ABC**, **Primakoff** and  **$^{57}\text{Fe}$**  axions (unconstrained components in the fit)
- ▶ Axion hypothesis **still favoured over  $B_0 + \text{tritium}$**  at  $2.0\sigma$



# Solar axion results





PRD 102 (2020) 072004



- ▶ 3D 90% C.L. volume  $g_{ae}$  vs.  $g_{ae} g_{a\gamma}$  vs.  $g_{ae} g_{an}^{eff}$

# Solar axion results

PRD 102 (2020) 072004




- ▶ 3D 90% C.L. volume  $g_{ae}$  vs.  $g_{ae}g_{ay}$  vs.  $g_{ae}g_{an}^{\text{eff}}$
- ▶ Exclusion of  $g_{ae} = 0$  OR  $g_{ae}g_{ay} = g_{ae}g_{an}^{\text{eff}} = 0$
- ▶ **Strong tension** with stellar cooling constraints

# The $\nu$ magnetic moment hypothesis

PRD 102 (2020) 072004



- ▶ SM extensions → **magnetic moment  $\mu_\nu$**  enhancing  $\nu$ -e<sup>-</sup> cross section **at low energy**
- ▶ Cherry on top,  $\mu_\nu \gtrsim 10^{-15} \mu_B$  → neutrinos may be **Majorana** fermions



# The $\nu$ magnetic moment hypothesis

PRD 102 (2020) 072004

- SM extensions → **magnetic moment  $\mu_\nu$**  enhancing  $\nu$ -e<sup>-</sup> cross section **at low energy**
- Cherry on top,  $\mu_\nu \gtrsim 10^{-15} \mu_B$  → neutrinos may be **Majorana** fermions





# The $\nu$ magnetic moment hypothesis

PRD 102 (2020) 072004

- SM extensions → **magnetic moment  $\mu_\nu$**  enhancing  $\nu$ -e<sup>-</sup> cross section **at low energy**
- Cherry on top,  $\mu_\nu \gtrsim 10^{-15} \mu_B$  → neutrinos may be **Majorana** fermions
- Strong tension** with astrophysical limits but consistent with comparable Borexino results


Neutrino magnetic  
moment hypothesis  
favoured over  $B_0$  at  **$3.2\sigma$**



# The bosonic Dark Matter hypothesis

PRD 102 (2020) 072004


- ▶ Searching for two main components as **monoenergetic peaks**
  - **Axion-Like Particles**  $\approx$  axions ( $\propto (g_{ae})^2$ ) with higher masses
  - **Dark photons**, could couple with photons ( $\propto \kappa^2$ ) and be absorbed by photoelectric effect





# The bosonic Dark Matter hypothesis

PRD 102 (2020) 072004

- ▶ Searching for two main components as **monoenergetic peaks**
  - **Axion-Like Particles**  $\approx$  axions ( $\propto (g_{ae})^2$ ) with higher masses
  - **Dark photons**, could couple with photons ( $\propto \kappa^2$ ) and be absorbed by photoelectric effect






- ▶ Most significant at  
 **$(2.3 \pm 0.2)$  keV/c<sup>2</sup>**  
(favoured at  $3\sigma$ )



# The bosonic Dark Matter hypothesis


PRD 102 (2020) 072004

- ▶ Searching for two main components as **monoenergetic peaks**
  - **Axion-Like Particles**  $\approx$  axions ( $\propto (g_{ae})^2$ ) with higher masses
  - **Dark photons**, could couple with photons ( $\propto \kappa^2$ ) and be absorbed by photoelectric effect
- ▶ Most significant at  **$(2.3 \pm 0.2)$  keV/c<sup>2</sup>** (favoured at  $3\sigma$ )
- ▶ No excess above  $3\sigma \rightarrow$  upper limits on  $g_{ae}$  and  $\kappa$
- ▶ **Best limits overall**



# The low-energy excess


PRD 102 (2020) 072004



Play with our data → <https://doi.org/10.5281/zenodo.4273099>

# The low-energy excess

PRD 102 (2020) 072004




Play with our data → <https://doi.org/10.5281/zenodo.4273099>



**What's next?**

# Meet XENONnT



# Meet XENONnT





# Meet XENONnT





# Meet XENONnT




# Meet XENONnT




# Meet XENONnT



# Meet XENONnT




# XENONnT take-home



@XENOExperiment

Data taking  
**Starting late 2020**



Active volume  
 $\times 3$

Background  
**1/6**

Axion–tritium  
discrimination  
**A few  
months**



# Thank you for your attention!

Dr. Erwann Masson

IJCLab – Orsay, France

✉ masson@lal.in2p3.fr

🐦 @DrErwannMasson



université  
PARIS-SACLAY

Xe  
XENON  
Dark Matter Project